



We ~~claim~~ claim:

1. A method of avoiding hyperphosphatemia while treating a patient having a kidney disorder comprising administering to said patient a vitamin D compound that has a minimal effect on serum phosphorus of said patient.
2. The method of claim 1 wherein said kidney disorder is uremia.
3. The method of claim 1 wherein said kidney disorder is renal failure.
4. The method of claim 1 wherein said vitamin D compound is administered together with a pharmaceutically acceptable excipient.
5. The method of claim 1 wherein said vitamin D compound is in a solid or liquid vehicle ingestible by and non-toxic to the patient.
6. The method of claim 1 wherein said vitamin D compound is a 19-nor-vitamin D compound.
7. The method of claim 6 wherein said 19-nor-vitamin D compound has the formula:




where  $X^1$  and  $X^2$  each represent, independently, hydrogen or a hydroxy-protecting group, and where R is represented by the structure below:



20

where the stereochemical center may have the R or S configuration, and where Z is selected from Y, -OY, -CH<sub>2</sub>OY, -C ≡ CY and -CH = CHY, where the double bond may have the cis or trans geometry, and where Y is selected from hydrogen, methyl, -CR<sup>5</sup>O and a radical of the structure.

25



30

where m and n, independently, represent integers from 0 to 5, where R<sup>1</sup> is selected from hydrogen, hydroxy, protected hydroxy, fluoro, trifluoromethyl, and C<sub>1-5</sub>- alkyl, which may be straight chain or branched and, optionally, bear a hydroxy or protected-hydroxy substituent, and where each of R<sup>2</sup>, R<sup>3</sup>, and R<sup>4</sup>, independently, is selected from hydrogen, fluoro, trifluoromethyl and C<sub>1-5</sub> alkyl, which may be straight-chain or branched, and optionally, bear a hydroxy or protected-hydroxy substituent, and where R<sup>1</sup> and R<sup>2</sup>, taken together represent an oxo group, or an alkylidene group, =CR<sup>2</sup>R<sup>3</sup>, or the group -(CH<sub>2</sub>)<sub>p</sub>-, where p is an integer from 2 to 5, and where R<sup>3</sup> and R<sup>4</sup>, taken together, represent an oxo group, or a group -(CH<sub>2</sub>)<sub>q</sub>-, where q is an integer from 2 to 5, where R<sup>5</sup> represents hydrogen, hydroxy, protected hydroxy, or C<sub>1-5</sub> alkyl, and where any of the groups at positions 20, 22 and 23, respectively in the side chain may be replaced by an oxygen atom.

35

40

45

8. The method of claim 1 where the said vitamin D compound is 1 $\alpha$ ,25-dihydroxy-19-nor-vitamin D<sub>3</sub>.

9. The method of claim 1 where the said vitamin D compound is 1 $\alpha$ -hydroxy-19-nor-vitamin D<sub>3</sub>.

10. The method of claim 1 where the said vitamin D compound is 1 $\alpha$ ,25-dihydroxy-19-nor-vitamin D<sub>2</sub>.

11. The method of claim 1 where the said vitamin D compound is 1 $\alpha$ -hydroxy-19-nor-vitamin D<sub>2</sub>.

12. The method of claim 1 where the said vitamin D compound is 1 $\alpha$ -hydroxy-19-nor-24-epi-vitamin D<sub>2</sub>.

13. The method of claim 1 where the said vitamin D compound is 1 $\alpha$ ,25-dihydroxy-19-nor-24-epi-vitamin D<sub>2</sub>.

14. The method of claim 1 where the said vitamin D compound is administered orally.

15. The method of claim 1 where the said vitamin D compound is administered parenterally.

16. The method of claim 1 where the said vitamin D compound is administered topically.

17. The method of claim 1 where the said vitamin D compound is administered in an amount from 1  $\mu$ g to about 500  $\mu$ g per day to the patient.

Add  
A27